

Occult Contralateral Central Lymph Node Metastases In

Tumor size >1 cm, the presence of benign nodule, extrathyroidal extension, central lymph node metastasis, lateral lymph node metastasis are independent predictors of contralateral occult carcinoma. Contralateral occult carcinoma is the independent predictor of recurrence. A 10-point risk-scoring model was established to predict the

**...
In high-risk patients with occult contralateral central LN metastases, as shown in our data, comprehensive ND including bilateral central neck compartment can help in preventing tumor recurrence in the contralateral lateral neck, although the prognostic significance of occult contralateral central LN metastases might be unknown because of the deficiency in long-term follow-up results. Prediction of ipsilateral and contralateral central lymph node metastasis in unilateral papillary thyroid carcinoma: a retrospective study. Qiang Chen, Xiu-He Zou, ... The incidence of occult nodal metastasis of PTC in central compartment has been reported to vary between 30% and 90% .
Occult Contralateral Central Lymph Node Metastases In
Occult Contralateral Central Lymph Node**

In high-risk patients with occult contralateral central LN metastases, as shown in our data, comprehensive ND including bilateral central neck compartment can help in preventing tumor recurrence in the contralateral lateral neck, although the prognostic significance of occult contralateral central LN metastases might be unknown because of the deficiency in long-term follow-up results.

**Occult Contralateral Central Lymph Node Metastases in ...
Occult LNM in the contralateral lateral neck was found in 36.5% of patients. Five or more ipsilateral central LNM may aid in predicting contralateral lateral LNM, and high-risk patients may be more at risk. The clinical benefit of prophylactic contralateral lateral ND remains doubtful, however.**

**Occult Contralateral Lateral Lymph Node Metastases in ...
Prediction of ipsilateral and contralateral central lymph node metastasis in unilateral papillary thyroid carcinoma: a retrospective study. Qiang Chen, Xiu-He Zou, ... The**

incidence of occult nodal metastasis of PTC in central compartment has been reported to vary between 30% and 90% .

Prediction of ipsilateral and contralateral central lymph

...

Occult Contralateral Central Lymph Node Metastases In
Author: s2.kora.com-2020-10-16T00:00:00+00:01 Subject:
Occult Contralateral Central Lymph Node Metastases In
Keywords: occult, contralateral, central, lymph, node,
metastases, in Created Date: 10/16/2020 2:36:26 PM

Occult Contralateral Central Lymph Node Metastases In
The relationships between occult LN metastases to the
contralateral central neck compartment and preoperative
image findings and clinicopathologic factors were analyzed.
RESULTS: Central neck LN metastases were present in 82.9%,
in which 34.3% had bilateral central neck involvement and
48.6% had unilateral ipsilateral central neck involvement.

Occult contralateral central lymph node metastases in ...
Occult Contralateral Central Lymph Node Occult LNM in the
contralateral lateral neck was found in 36.5% of patients.
Five or more ipsilateral central LNM may aid in predicting
contralateral lateral LNM, and high-risk patients may be
more at risk. The clinical benefit of prophylactic
contralateral lateral ND remains doubtful, however.

Occult Contralateral Central Lymph Node Metastases In
Therapeutic lateral neck dissection (ND) is recommended for
N1b papillary thyroid carcinoma (PTC), while prophylactic
contralateral lateral ND is not. Given the paucity of data,
we investigated the frequency of and risk factors for
occult lymph node metastases (LNM) in the contralateral
lateral neck for N1b patients. This is a retrospective
study conducted at a cancer center.

Occult Contralateral Lateral Lymph Node Metastases in ...
In our study, lymph node metastases were found in 18.7% (72
in 386 cases) patients with contralateral occult PTC
(P=0.004). Multivariate analysis showed that lymph node
metastasis was an independent predictor of contralateral
occult PTC for patients whose pre-operation US showed a
unilateral PTC.

Predictive factors of contralateral occult carcinoma in ...
The frequency and predictive factors for occult contralateral carcinoma were analyzed with respect to the following variables: age, gender, family history, tumor size, presence of Hashimoto thyroiditis, extrathyroidal extension, central lymph node metastasis, multifocality of primary tumor, or BRAFT1799A mutation.

The BRAFT1799A mutation is not associated with occult ... The frequency, pattern, and predictive factors for occult contralateral carcinoma in these patients were analyzed with respect to the following variables: age, gender, tumor size, multifocality of primary tumor, presence of perithyroidal invasion, lymphovascular invasion or capsular invasion, presence of central lymph node metastasis, and the ...

Occult Contralateral Carcinoma in Patients with Unilateral ...
all enlarged nodes (size $\geq 5\text{mm}$) by EBUS. However, procedure-time is limited by patient comfort when moderate sedation is used. It is unclear if EBUS-staging should start with hilar N3 nodes or whether starting with mediastinal N3 nodes suffices. Knowing the probability of PET-CT negative hilar N3 nodes harboring occult metastasis can inform this

Is Biopsy of Contralateral Hilar N3 Lymph Nodes that Are ...
However, Koo et al. insisted on bilateral central neck dissection due to the relatively high rate of occult contralateral central lymph node metastasis (34.3%), especially in multifocal primary ...

Occult Contralateral Central Lymph Node Metastases in ...
Objective To investigate the prevalence and distribution of cervical lymph node metastasis (LNM) in locally advanced supraglottic squamous cell carcinoma (LASCC) and guide the delineation of clinical lymph node target volumes. Materials and Methods We reviewed patients defined as LASCC from January 2000 to December 2017 in our hospital. The primary tumor was operated on using partial or total ...

Frontiers | Patterns of Cervical Lymph Node Metastasis in

...

Tumor size >1 cm, the presence of benign nodule, extrathyroidal extension, central lymph node metastasis, lateral lymph node metastasis are independent predictors of contralateral occult carcinoma. Contralateral occult carcinoma is the independent predictor of recurrence. A 10-point risk-scoring model was established to predict the

...

Frontiers | Management of Clinically Solitary Papillary ...

Results: Occult contralateral lateral LNM were found in 23/63 patients (36.5%) and were significantly associated with the number of metastatic nodes in the central compartment ipsilateral to the tumor ($p=.02$) and with microscopic tumor foci in the contralateral thyroid lobe ($p=0,017$).

OP72 – Predictive factors for occult contralateral lateral

...

contralateral nodes. FAQ. Medical Information Search ... Lymph Nodes Sinoatrial Node Atrioventricular Node Ranvier's Nodes Motor Cortex Axilla Forelimb Efferent Pathways Somatosensory Cortex Neural Pathways Olivary Nucleus Hindlimb Pyramidal ... Central Stomach Neoplasms Ureteral Obstruction Testicular Neoplasms Carotid Artery Diseases Ischemia ...

Contralateral nodes. Medical search. FAQ

Background: Prophylactic central lymph node dissection (CLND) in patients with papillary thyroid carcinoma (PTC) remains controversial and predictive factors for central lymph node (CLN) metastasis in unilateral PTC cases are not well defined. The aims of this study were to evaluate the rate of ipsilateral and contralateral CLN metastasis and to determine the clinicopathologic factors ...

Prediction of ipsilateral and contralateral central lymph

...

BOSTON – More than a third of 63 patients with unilateral papillary thyroid carcinoma and ipsilateral lymph node metastases also had occult, contralateral, lateral-neck lymph nodes with metastases, but the low reported rate of

contralateral neck recurrence has raised the question of whether routine resection of these contralateral lymph nodes benefits patients.

**Contralateral nodal thyroid metastases show slow ...
<https://jes-online.org> 191 ABSTRACT Purpose: The purpose of this study was to assess the presence of contralateral occult malignant foci (OMF) among patients who confined unilateral papillary thyroid carcinomas (PTCs). Methods: From January 2010 to December 2012, we retrospectively evaluated 714 patients who received total thyroidectomy with central lymph node (LN) dissection after being**

**Original Article Prediction of Contralateral Occult ...
Predictive factors for ipsilateral or contralateral central lymph node metastasis in unilateral papillary thyroid carcinoma. Ann Surg. 2009;249:840-4 18. Lee CC, Lin YS, Kang BH, Chang KP, Chi CC, Lin MY. et al. Incorporation of log odds of positive lymph nodes into the AJCC TNM classification improves prediction of survival in oral cancer.**

Original Article Prediction of Contralateral Occult ...

Contralateral nodes. Medical search. FAQ

contralateral nodes. FAQ. Medical Information Search ... Lymph Nodes Sinoatrial Node Atrioventricular Node Ranvier's Nodes Motor Cortex Axilla Forelimb Efferent Pathways Somatosensory Cortex Neural Pathways Olivary Nucleus Hindlimb Pyramidal ... Central Stomach Neoplasms Ureteral Obstruction Testicular Neoplasms Carotid Artery Diseases Ischemia ...

Occult Contralateral Central Lymph Node Metastases In Author:

s2.kora.com-2020-10-16T00:00:00+00:01 Subject: Occult Contralateral Central Lymph

Metastases In Keywords: occult, contralateral, central, lymph, node, metastases, in Cre

Date: 10/16/2020 2:36:26 PM

Background: Prophylactic central lymph node dissection (CLND) in patients with papillary thyroid carcinoma (PTC) remains controversial and predictive factors for central lymph node (CLN) metastasis in unilateral PTC cases are not well defined. The aims of this study were to evaluate the rate of ipsilateral and contralateral CLN metastasis and to determine the clinicopathologic factors ...

Prediction of ipsilateral and contralateral central lymph ...

Occult Contralateral Lateral Lymph Node Metastases in ...

The frequency, pattern, and predictive factors for occult contralateral

carcinoma in these patients were analyzed with respect to the following variables: age, gender, tumor size, multifocality of primary tumor, presence of perithyroidal invasion, lymphovascular invasion or capsular invasion, presence of central lymph node metastasis, and the ...

The frequency and predictive factors for occult contralateral carcinoma were analyzed with respect to the following variables: age, gender, family history, tumor size, presence of Hashimoto thyroiditis, extrathyroidal extension, central lymph node metastasis, multifocality of primary tumor, or BRAFT1799A mutation.

Occult Contralateral Central Lymph Node Occult LNM in the contralateral lateral neck was found in 36.5% of patients. Five or more ipsilateral central LNM may aid in predicting contralateral lateral LNM, and high-risk patients may be more at risk. The clinical benefit of prophylactic contralateral lateral ND remains doubtful, however.

The BRAFT1799A mutation is not associated with occult ... Therapeutic lateral neck dissection (ND) is recommended for N1b papillary thyroid carcinoma (PTC), while prophylactic contralateral lateral ND is not. Given the paucity of data, we investigated the frequency of and risk factors for occult lymph node metastases (LNM) in the contralateral lateral neck for N1b patients. This is a retrospective study conducted at a cancer center.

The relationships between occult LN metastases to the contralateral central neck compartment and preoperative image findings and clinicopathologic factors were analyzed. RESULTS: Central neck LN metastases were present in 82.9%, in which 34.3% had bilateral central neck involvement and 48.6% had unilateral ipsilateral central neck involvement.

OP72 – Predictive factors for occult contralateral lateral ...

Objective To investigate the prevalence and distribution of cervical lymph node metastasis (LNM) in locally advanced supraglottic squamous cell carcinoma (LASCC) and guide the delineation of clinical lymph node target volumes. Materials and Methods We reviewed patients defined as LASCC from January 2000 to December 2017 in our hospital. The primary tumor was operated on using partial or total ...

Contralateral nodal thyroid metastases show slow ...

Occult Contralateral Carcinoma in Patients with Unilateral ...

Occult LNM in the contralateral lateral neck was found in 36.5% of patients. Five or more ipsilateral central LNM may aid in predicting contralateral lateral LNM,

and high-risk patients may be more at risk. The clinical benefit of prophylactic contralateral lateral ND remains doubtful, however.

Occult Contralateral Central Lymph Node

Is Biopsy of Contralateral Hilar N3 Lymph Nodes that Are ...

Occult Contralateral Central Lymph Node Metastases in ...

Results: Occult contralateral lateral LNM were found in 23/63 patients (36.5%) and were significantly associated with the number of metastatic nodes in the central compartment ipsilateral to the tumor ($p=.02$) and with microscopic tumor foci in the contralateral thyroid lobe ($p=0,017$).

Frontiers | Management of Clinically Solitary Papillary ...

BOSTON – More than a third of 63 patients with unilateral papillary thyroid carcinoma and ipsilateral lymph node metastases also had occult, contralateral, lateral-neck lymph nodes with metastases, but the low reported rate of contralateral neck recurrence has raised the question of whether routine resection of these contralateral lymph nodes benefits patients.

Predictive factors of contralateral occult carcinoma in ...

<https://jes-online.org> 191 ABSTRACT Purpose: The purpose of this study was to assess the presence of contralateral occult malignant foci (OMF) among patients who confined unilateral papillary thyroid carcinomas (PTCs). Methods: From January 2010 to December 2012, we retrospectively evaluated 714 patients who received total thyroidectomy with central lymph node (LN) dissection after being

Occult contralateral central lymph node metastases in ...

Frontiers | Patterns of Cervical Lymph Node Metastasis in ...

In our study, lymph node metastases were found in 18.7% (72 in 386 cases) patients with contralateral occult PTC ($P=0.004$). Multivariate analysis showed that lymph node metastasis an independent predictor of contralateral occult PTC for patients whose pre-operation US showed a unilateral PTC.

Occult Contralateral Central Lymph Node

In high-risk patients with occult contralateral central LN metastases, as shown in our data, comprehensive ND including bilateral central neck compartment can help in preventing tumor recurrence in the contralateral lateral neck, although the prognostic significance of occult contralateral central LN metastases might be unknown because of the deficiency in long-term follow-up results.

Occult Contralateral Central Lymph Node Metastases in ...

Occult LNM in the contralateral lateral neck was found in 36.5% of patients. Five or more ipsilateral central LNM may aid in predicting contralateral lateral LNM, and high-risk patients may be more at risk. The clinical benefit of prophylactic contralateral lateral ND remains doubtful, however.

Occult Contralateral Lateral Lymph Node Metastases in ...

Prediction of ipsilateral and contralateral central lymph node metastasis in unilateral papillary

thyroid carcinoma: a retrospective study. Qiang Chen, Xiu-He Zou, ... The incidence of occult nodal metastasis of PTC in central compartment has been reported to vary between 30% and .

Prediction of ipsilateral and contralateral central lymph ...

Occult Contralateral Central Lymph Node Metastases In Author:

s2.kora.com-2020-10-16T00:00:00+00:01 Subject: Occult Contralateral Central Lymph Node

Metastases In Keywords: occult, contralateral, central, lymph, node, metastases, in Created D

10/16/2020 2:36:26 PM

Occult Contralateral Central Lymph Node Metastases In

The relationships between occult LN metastases to the contralateral central neck compartment and preoperative image findings and clinicopathologic factors were analyzed. RESULTS: Central neck LN metastases were present in 82.9%, in which 34.3% had bilateral central neck involvement and 48.6% had unilateral ipsilateral central neck involvement.

Occult contralateral central lymph node metastases in ...

Occult Contralateral Central Lymph Node Occult LNM in the contralateral lateral neck was found in 36.5% of patients. Five or more ipsilateral central LNM may aid in predicting contralateral lateral LNM, and high-risk patients may be more at risk. The clinical benefit of prophylactic contralateral lateral ND remains doubtful, however.

Occult Contralateral Central Lymph Node Metastases In

Therapeutic lateral neck dissection (ND) is recommended for N1b papillary thyroid carcinoma (PTC), while prophylactic contralateral lateral ND is not. Given the paucity of data, we investigated the frequency of and risk factors for occult lymph node metastases (LNM) in the contralateral lateral neck for N1b patients. This is a retrospective study conducted at a cancer center.

Occult Contralateral Lateral Lymph Node Metastases in ...

In our study, lymph node metastases were found in 18.7% (72 in 386 cases) patients with contralateral occult PTC ($P=0.004$). Multivariate analysis showed that lymph node metastasis is an independent predictor of contralateral occult PTC for patients whose pre-operation US showed a unilateral PTC.

Predictive factors of contralateral occult carcinoma in ...

The frequency and predictive factors for occult contralateral carcinoma were analyzed with respect to the following variables: age, gender, family history, tumor size, presence of Hashimoto thyroiditis, extrathyroidal extension, central lymph node metastasis, multifocality of primary tumor, or BRAFT1799A mutation.

The BRAFT1799A mutation is not associated with occult ...

The frequency, pattern, and predictive factors for occult contralateral carcinoma in these patients were analyzed with respect to the following variables: age, gender, tumor size, multifocality of primary tumor, presence of perithyroidal invasion, lymphovascular invasion or capsular invasion, presence of central lymph node metastasis, and the ...

Occult Contralateral Carcinoma in Patients with Unilateral ...

all enlarged nodes (size \geq 5mm) by EBUS. However, procedure-time is limited by patient comfort when moderate sedation is used. It is unclear if EBUS-staging should start with hilar N3 nodes

whether starting with mediastinal N3 nodes suffices. Knowing the probability of PET-CT negative hilar N3 nodes harboring occult metastasis can inform this

Is Biopsy of Contralateral Hilar N3 Lymph Nodes that Are ...

However, Koo et al. insisted on bilateral central neck dissection due to the relatively high rate of occult contralateral central lymph node metastasis (34.3%), especially in multifocal primary ...

Occult Contralateral Central Lymph Node Metastases in ...

ObjectiveTo investigate the prevalence and distribution of cervical lymph node metastasis (LNM) in locally advanced supraglottic squamous cell carcinoma (LASCC) and guide the delineation of clinical lymph node target volumes.
Materials and MethodsWe reviewed patients defined as LASCC from January 2000 to December 2017 in our hospital. The primary tumor was operated on using partial or total ...

Frontiers | Patterns of Cervical Lymph Node Metastasis in ...

Tumor size >1 cm, the presence of benign nodule, extrathyroidal extension, central lymph node metastasis, lateral lymph node metastasis are independent predictors of contralateral occult carcinoma. Contralateral occult carcinoma is the independent predictor of recurrence. A 10-point risk-scoring model was established to predict the ...

Frontiers | Management of Clinically Solitary Papillary ...

Results: Occult contralateral lateral LNM were found in 23/63 patients (36.5%) and were significantly associated with the number of metastatic nodes in the central compartment ipsilateral to the tumor ($p=0.02$) and with microscopic tumor foci in the contralateral thyroid lobe ($p=0.017$).

OP72 – Predictive factors for occult contralateral lateral ...

contralateral nodes. **FAQ.** Medical Information Search ... Lymph Nodes Sinoatrial Node Atrioventricular Node Ranvier's Nodes Motor Cortex Axilla Forelimb Efferent Pathways Somatosensory Cortex Neural Pathways Olivary Nucleus Hindlimb Pyramidal ... Central Stomach Neoplasms Ureteral Obstruction Testicular Neoplasms Carotid Artery Diseases Ischemia ...

Contralateral nodes. Medical search. **FAQ**

Background: Prophylactic central lymph node dissection (CLND) in patients with papillary thyroid carcinoma (PTC) remains controversial and predictive factors for central lymph node (CLN) metastasis in unilateral PTC cases are not well defined. The aims of this study were to evaluate the rate of ipsilateral and contralateral CLN metastasis and to determine the clinicopathologic factors ...

Prediction of ipsilateral and contralateral central lymph ...

BOSTON – More than a third of 63 patients with unilateral papillary thyroid carcinoma and ipsilateral lymph node metastases also had occult, contralateral, lateral-neck lymph nodes with metastases, but the low reported rate of contralateral neck recurrence has raised the question whether routine resection of these contralateral lymph nodes benefits patients.

Contralateral nodal thyroid metastases show slow ...

<https://jes-online.org> 191 **ABSTRACT** Purpose: The purpose of this study was to assess the presence of contralateral occult malignant foci (OMF) among patients who confined unilateral papillary thyroid carcinomas (PTCs). Methods: From January 2010 to December 2012, we retrospectively evaluated 714 patients who received total thyroidectomy with central lymph node ...

(LN) dissection after being

Original Article Prediction of Contralateral Occult ...

Predictive factors for ipsilateral or contralateral central lymph node metastasis in unilateral papillary thyroid carcinoma. *Ann Surg.* 2009;249:840-4 18. Lee CC, Lin YS, Kang BH, Chang KP, Chi CC, Lin MY. et al. Incorporation of log odds of positive lymph nodes into the AJCC TNM classification improves prediction of survival in oral cancer.

Predictive factors for ipsilateral or contralateral central lymph node metastasis in unilateral papillary thyroid carcinoma. *Ann Surg.* 2009;249:840-4 18. Lee CC, Lin YS, Kang BH, Chang KP, Chi CC, Lin MY. et al. Incorporation of log odds of positive lymph nodes into the AJCC TNM classification improves prediction of survival in oral cancer.

all enlarged nodes (size \geq 5mm) by EBUS. However, procedure-time is limited by patient comfort when moderate sedation is used. It is unclear if EBUS-staging should start with hilar N3 nodes or whether starting with mediastinal N3 nodes suffices. Knowing the probability of PET-CT negative hilar N3 nodes harboring occult metastasis can inform this

However, Koo et al. insisted on bilateral central neck dissection due to the relatively high rate of occult contralateral central lymph node metastasis (34.3%), especially in multifocal primary ...